

## **Sleep Disturbance and Daytime Sleepiness in Patients with Cirrhosis: a Case Control Study**

Barbara Mostacci<sup>1</sup>, Monica Ferlisi<sup>1</sup>, Alessandro Baldi Antognini<sup>2</sup>, Claudia Sama<sup>3</sup>, Cristina Morelli<sup>3</sup>, Susanna Mondini<sup>1</sup>, Fabio Cirignotta<sup>1</sup>,

*<sup>1</sup>Unit of Neurology, <sup>3</sup>Dept of Internal Medicine , S.Orsola-Malpighi Hospital, University of Bologna, Italy <sup>2</sup> Dept of Statistical Sciences, University of Bologna, Italy*

Author 's address : prof. Fabio Cirignotta UO Neurologia , Policlinico S.Orsola-Malpighi , via Albertoni 15 , 40138 Bologna tel. +39 051 6262589 fax +39 05163562640 e-mail [fabio.cirignotta@aosp.bo.it](mailto:fabio.cirignotta@aosp.bo.it)

---

<sup>°</sup> Correspondence: Prof. Fabio Cirignotta, UO Neurologia, Policlinico S. Orsola-Malpighi, Via Albertoni 15, 40138 Bologna, Italy. Email: [fabio.cirignotta@aosp.bo.it](mailto:fabio.cirignotta@aosp.bo.it) Fax: (0039)0516362640

**Abstract:**

**Objective:** Sleep disturbance and excessive daytime sleepiness have been reported in patients with hepatic cirrhosis. The objective of this study was to evaluate daytime somnolence and sleep complaints in a group of 178 patients with cirrhosis compared to a control group.

**Method:** Sleep features and excessive daytime sleepiness were evaluated by the Basic Nordic Sleep Questionnaire (BNSQ) and the Epworth Sleepiness Scale (ESS). We collected clinical and laboratory data, neurological assessment and EEG recording in cirrhotic patients.

**Results:** Patients with cirrhosis complained of more daytime sleepiness ( $p < 0.005$ ), sleeping badly at least three times a week ( $p < 0.005$ ), difficulties falling asleep ( $p < 0.01$ ) and frequent nocturnal awakening ( $p < 0.005$ ) than controls. We found a poor correlation between sleep disorders and clinical or laboratory parameters. **Conclusion:** Our results confirm previous literature reports suggesting a high prevalence of sleep disturbance in patients with cirrhosis. Insomnia and daytime sleepiness are the main complaints. Sleep disorders are probably a multifactorial phenomenon.

**Key words:** liver cirrhosis, insomnia, daytime somnolence.

## **Introduction**

Sleep disturbance and excessive daytime somnolence are common in patients with cirrhosis [1, 2, 3, 4]. In addition, a disturbance of sleep is recognized as one of the early signs of hepatic encephalopathy [5, 6, 7]. Reversal of sleep rhythm, drowsiness and lethargy are classic signs of this disease, and their presence and entity are used to define the clinical stages of hepatic encephalopathy [8, 9, 10, 11].

The most common feature of the sleep pattern in patients with cirrhosis is fragmented nocturnal sleep caused by frequent nocturnal awakenings, and a daytime functioning affected by frequent episodes of undesired sleepiness and more prolonged napping time [1, 2]. The sleep-wake cycle was also shifted with activity toward the later hours of the day. This was ascribed to the displacement toward later hours in the 24-hour profile of plasma melatonin, common in cirrhosis [12]. However, the existence of a phase delay in cirrhotic patients and its relation with the melatonin rhythm remains controversial [13]. In addition, few data showed a correlation between sleep impairment and the clinical parameters of liver disease. There are limited data on the true prevalence of sleep disturbance in cirrhotic patients compared with a control group.

For these reasons, we conducted a case control study to assess the prevalence and characteristics of sleep disturbance and excessive daytime sleepiness in patients with cirrhosis compared with a sex and age matched non-cirrhotic population, and to determine their correlation with clinical parameters.

## **Materials and methods**

### **Participants**

*Patients with cirrhosis.* We studied a consecutive series of 178 patients with hepatic cirrhosis. All patients with a diagnosis of liver cirrhosis not alcohol related seen at our Neurology Unit were invited to participate in this study. Patients were referred to our Unit either for neurological screening for orthotopic liver transplantation (112 patients) or for any neurological complaint (including hepatic encephalopathy). Patients with post-alcoholic cirrhosis were excluded due to the possible independent influence of ethanol on the central nervous system and quality of sleep.

One hundred and fifteen males and sixty-three females, whose mean age was 52.12 years (ranging from 17 to 71 years) were enrolled.

The diagnosis of cirrhosis was based on a compatible clinical history, radiological studies, and liver biopsy when available.

At the time of evaluation, 23.5% were Child-Pugh class A patients, 48.2% class B and 28.3% class C. The aetiology of cirrhosis was viral hepatitis in 78.6% (HBV 18.5%; HCV 53.4%; HBV+HCV 6.7%), autoimmune disease in 9% and diverse causes in 11.4%. 24.2% of patients referred past episodes of hepatic encephalopathy, 6.2% had current minimal hepatic encephalopathy and 68% had never had episodes of encephalopathy. 136 patients were on the list for orthotopic liver transplantation.

*Healthy controls.* A control group was selected from healthy subjects, mostly blood donors at our hospital and employees from our institution. The group included 178 individuals sex and age matched with the cirrhotic patients (+/- 1 year). Mean age of normal controls was 52.06 years (ranging from 17 to 72 years).

The study was conducted according to the principles established in Helsinki and approved by the local ethical committee.

## **Procedures**

*Neurological assessment.* A complete neurological assessment was performed in all patients with cirrhosis. One-hundred-thirteen patients underwent EEG recording.

*Sleep Evaluation.* All patients with cirrhosis were assessed by a sleep specialist with a semi-structured 20 minute interview on their sleep habits and complaints. All subjects were administered two standardised questionnaires: the Basic Nordic Sleep Questionnaire (BNSQ) and the Epworth Sleepiness Scale (ESS).

The BNSQ comprises 22 questions investigating sleep habits and complaints of insomnia, difficulty falling asleep, night awakenings, early morning awakening, use of hypnotic drugs, napping habits, daytime sleepiness, snoring and sleep apnoea [14].

We added two questions to better define the severity of daytime sleepiness and the circumstance in which sleepiness occurred. The questions added were: “Do you feel excessively sleepy during the daytime?” 1-never 2-sometimes after lunch 3-often or always after lunch 4-even at other times of the day 5-throughout the day; and “Sleepiness is” 1-absent 2-slight 3-moderate 4-severe 5-very severe. Most of the questions ask for the frequency of the symptom (never or less than once per month, less than once per week, on 1-2 days per week, on 3-5 days per week, daily or almost daily), other than the severity of the symptom.

The ESS is a standardised scale scoring the level of daytime sleepiness by rating the chance the subject would doze off in eight different real-life situations [15]. The questionnaires were administered by a physician.

All subjects were classified on the basis of their answers to the BNSQ questions as Sleepy (S) if they complained of excessive daytime sleepiness, at least moderate and not only in the postprandial hours at least three times a week, Postprandial Sleepy (PS) if they complained of

excessive daytime sleepiness at least moderate only after lunch at least three times a week, or Not Sleepy (NS) if they did not fit these conditions. Subjects were classified as Insomniacs if they reported sleeping badly or rather badly at least three times a week.

*Clinical and laboratory data.* We included in the database information on education, employment, routine biochemistry, including albumin and serum bilirubin, prothrombin activity, cholesterol, alkaline phosphatase, blood ammonia, transaminase,  $\gamma$ -glutamyl transpeptidase,  $\gamma$ -globulins, serum triglyceride level,  $\alpha$ -fetoprotein, blood nitrogen and current therapy. We also investigated the presence and severity of ascites, episodes of encephalopathy, gastrointestinal bleeding and jaundice.

### **Statistical Analysis**

We used t test, variance analysis (ANOVA), Wilcoxon and Mann-Whitney test. Data were analysed by the McNemar test. We related all questionnaire answers to blood ammonia value with Pearson's correlation.

### **Results**

Patients with cirrhosis showed a significantly higher prevalence of sleep disturbance than the healthy control group.

Twenty-six per cent of patients complained of sleeping badly or almost badly at least three times a week ("insomniacs"), whereas this complaint was infrequent in the healthy subjects (7.9%) ( $p < 0.005$ ).

Patients with cirrhosis complaining of "insomnia" reported more often both difficulty initiating and maintaining sleep but they also differed from "non-insomniac" patients with cirrhosis for a higher prevalence of daytime sleepiness ( $p < 0.001$ ) and more frequent naps

( $p < 0.01$ ). No difference was observed between groups regarding early morning awakening and snoring.

Patients with cirrhosis showed a significantly higher prevalence of daytime sleepiness ( $p < 0.005$ ): 18.5% were classified as Sleepy (vs. 3.9% of controls), 30.9% referred Postprandial Sleepiness (vs. 18.5% of controls) and 50.6% were classified as Not Sleepy (vs. 77.5% of controls).

Among patients with cirrhosis 55% referred habitual naps vs. 23% of healthy controls; the patients with cirrhosis reported more prolonged napping time. (table 1). Among cirrhotic patients, the Sleepy group showed worsening parameters of nocturnal sleep quality, like nocturnal awakening ( $p < 0.05$ ), difficulties falling asleep ( $p < 0.05$ ), complaint of sleeping badly or almost badly at least three times a week ( $p < 0.005$ ) and referred sleep apnea ( $p < 0.05$ ). Patients with cirrhosis had a mean ESS score of 6.66 (vs. 6.17 in healthy controls), and 15.7% of them had a score higher than 10 (vs. 12.9% in healthy controls), but the difference was not statistically significant. *The patients with the highest scores, although not pathological, were patients complaining of "insomnia"*. Among patients with cirrhosis those with current minimal hepatic encephalopathy were more frequently "sleepy" ( $p < 0.005$ ) and "insomniac" ( $p < 0.05$ ).

**Table 1** Characteristics of sleep among groups

No significant difference was observed between Sleepy and Postprandial or Not Sleepy cirrhotic patients with regard to Child-Pugh score, development of ascites, aetiology of cirrhosis, previous encephalopathy and routine biochemistry. The mean value of blood

ammonia was 61.79 mcg/dl in Postprandial and Not Sleepy, and 76.88 in Sleepy patients, but the difference was not significant (ANOVA). We tried to divide patients into two groups, according to pathological or normal blood ammonia value ( $\leq$  or  $>$  80 mcg/dl), and the two groups were compared according to parameters of sleep habits: they did not show significant differences. Moreover, blood ammonia value did not show any significant correlation with parameters of sleep quality.

Bedtime and wake-up time of patients with cirrhosis and healthy controls were similar (mean value of bedtime in cirrhotics: 23.16 vs. controls: 23.53; mean value of wake-up time in cirrhotics: 7.46 vs. controls: 7.36). Even comparing cirrhotic patients with sleep disturbance and those without, these parameters did not differ between groups (mean value of bedtime and wake-up time in insomniac cirrhotic group: 23.07 and 7.18; in non-insomniac cirrhotic group: 23.19 and 7.15).

## **Discussion**

Our study confirms that patients with cirrhosis frequently have sleep disorders. We found a significantly higher prevalence of parameters of poor sleep quality, like difficulties falling asleep, nocturnal awakenings and complaints of sleeping badly in patients with cirrhosis than in healthy controls. In addition, daytime functioning of these patients was affected by excessive sleepiness and more prolonged napping time. A higher prevalence of daytime sleepiness disclosed by the BNSQ questionnaire was not accompanied by a significant difference in the Epworth Sleepiness Scale (ESS). This may be because the ESS, validated in narcoleptics and OSAS patients, is not sufficiently accurate in estimating sleepiness in an inactive population like patients with cirrhosis [16]. As reported in previous studies, we found a poor correlation between clinical and laboratory parameters and sleep disturbance or daytime somnolence [17, 18]. We did not find a phase delay in the sleep patterns of patients



with cirrhosis, although the sleep questionnaire we used is not an accurate tool to identify circadian abnormalities.

Another limitation of our study is the difficulty defining and estimating sleepiness on the basis of subjective patient evaluation. The first problem is to distinguish sleepiness from common symptoms in cirrhosis, like fatigue, tiredness and lack of energy.

Sleepiness could also be linked to the limitations of a chronic disease forcing subjects to lead very sedentary lives, thereby increasing the risk of episodes of undesired sleepiness. In addition, the frequent nocturnal awakenings could be related to more episodes of nicturia because of diuretic treatment. In this regard, another limitation in our study is the lack of a control group with another chronic disease (e.g. chronic renal failure). Sleep difficulty in patients with cirrhosis could be related to a specific dysregulation of the histamine neurotransmitter system ( 19,20), and a comparison with other chronic diseases could yield information useful to estimate the real impact of hepatic failure on the sleep pattern and its pathogenesis.

Snoring and reported apnoeas did not significantly differ from healthy controls. However, the finding that snoring was more common in “sleepy” than in “post-prandial sleepy” and “not sleepy” patients with cirrhosis in our cohort could strengthen the hypothesis that somnolence in cirrhotic patients is favoured at least in part by an obstructive apnoea syndrome during sleep ( 21,22,23) .

## References

1. Cordoba J, Cabrera J, Lataif L, Penev P, Zee P, Blei A. T. High prevalence of sleep disturbance in cirrhosis. *Hepatology* 1998;27(2):339-345.
2. Bergonzi P, Bianco A, Mazza S, Mennuni G. Night sleep organization in patients with severe hepatic failure. *Eur Neurol* 1978;17:271-275.
3. Mechtcheriakov S, Graziadei IW, Kugener A, Wiedemann J, Galbavy C, Hinterhuber H, Marksteiner J, et al. Multidimensional assessment of neuro-psychiatric symptoms in patients with low-grade hepatic encephalopathy: a clinical rating scale. *World J Gastroenterol* 2005;11(37):5893-8.
4. Newton JL, Gibson GJ, Tomlinson M, Wilton K, Jones D. Fatigue in primary biliary cirrhosis is associated with excessive daytime somnolence. *Hepatology* 2006;44(1):91-8.
5. Velissaris D, Solomou E, Kalogeropoulos A, Georgiopolou V, Thomopoulos C, Karatza C. Sleep disorders and brain MRI as early indicators of subclinical hepatic encephalopathy. *Hepatogastroenterology* 2006;53(67):51-4.
6. Martino ME, Romero-Vives M, Fernandez-Lorente J, De Vicente E, Barcena R, Gaztelu JM. Sleep electroencephalogram alterations disclose initial stage of encephalopathy. *Methods Find Exp Clin Pharmacol* 2002;24 (Suppl):119-22.
7. Lewis M, Howdle PD. The neurology of liver failure. *Q J Med* 2003;96:623-633.
8. Jones EA, Weissenborn K. Neurology and the liver. *J Neurol Neurosurg Psychiatry* 1997;63:279-293.
9. Blei AT, Cordoba J, et al. Hepatic encephalopathy. *The American Journal of Gastroenterology* 2001;96(7): 1968-1975.
10. Sherlock S, Summerskill WHJ, White LP, Phear EA. Portal-systemic encephalopathy; neurological complications of liver disease. *The Lancet* 1954;267(6836):454-457.
11. Ortiz M, Jacas C, Cordoba J. Minimal hepatic encephalopathy: diagnosis, clinical significance and recommendations. *J Hepatol* 2005;42(Suppl 1):S45-53.
12. Steindl PE, Finn B, Bendok B, Rothke S, Zee PC, Blei AT. Disruption of the diurnal rhythm of plasma melatonin in cirrhosis. *Ann Internal Med* 1995;123:274-277.

13. Hourmand-Ollivier I, Piquet MA, Toudic JP, Denise P, Dao T. Actigraphy: A new diagnostic tool for hepatic encephalopathy. *World J Gastroenterol* 2006;12(14):2243-2244.
14. Partinem M, Gislason T. Basic North Sleep Questionnaire (BNSQ): a quantitated measure of subjective sleep complaints. *Sleep Res* 1995;4(S1):150-155.
15. Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep* 1991;14: 540-545.
16. Vignatelli L, Mattarozzi K, Zanatta C, Stracciari A. Cognitive function and Epworth Sleepiness Scale in minimal hepatic encephalopathy. *European Journal of Neurology* 2001;8(4):369.
17. Kurtz D, Zenglein JP, Imler M, Girardel M, Greenspan G, Peter B, Rohmer F. Etude du sommeil nocturne au cours de l'encephalopathie porto-cave. *Electroenceph Clin Neurophysiol* 1972 ;33:167-178.
18. Baldy-Moulinier M, Besset A, Calvet B, Michel H. Etude polygraphique nycthemerale des endormissement et des reveils au cours des encephalopathies hepatiques. *Rev Electroenceph Neurophysiol Clin* 1981;11:123-132.
- 19 Lozeva V, Tuomisto L, Tarhanen J, Butterworth RF .Increased concentrations of histamine and its metabolite, telemethylamine and down-regulation of histamine H3 receptor sites in autopsied brain tissue from cirrhotic patients who died in hepatic coma . *J Hepatol* 2003, 39(4) :522-7
- 20 Spahr L, Coeytaux MD, Giostra E, Hadengue A, Annoni JM . Histamine H1 blocker hydroxyzine improves sleep in patients with cirrhosis and minimal hepatic encephalopathy : a randomized controlled pilot trial. *Am J Gastroenterol* 2007, 102:744-753 ).
- 21 . Crespo J, Cifrian J, Pinto JA, Jimenez –Gomez A, Pons Romero F. Sleep apnea obstructive syndrome : a new complication previously underscribed in cirrhotic patients with ascites . *Am J Gastroenterol* 2003 , 98 : 2815-16
22. Ogata T, Nomura M, Nakaya Y, S.Ito . Evaluation of episodes of sleep apnea in patients with liver cirrhosis . *The journal of medical investigation* 2006 , 53 : 159-166
23. Nikaina I, Pastaka C, Zachou K, Dalekos GN, Gourgoulis K. Sleep apnea syndrome and early stage cirrhosis : a pilot study . *Eur J Gastroenterol Hepatol* 2006, 18 (1) :31-5

**Table 1** Characteristics of sleep among groups

	Cirrhotic Patients	Healthy Controls	P
Sleepy (%)	18.5	3.9	0.005
Post-prandial Sleepy (%)	30.9	18.5	
Not Sleepy (%)	50.6	77.5	
“Insomnias “(%)	26	7.9	0.005
Difficulty falling asleep (%)	21	8.5	0.01
Night Awakening (%)	71	43	0.005
Early awakening (%)	15	12	n.s.
Hypnotics (%)	4.5	2.3	n.s.
Morning sleepiness (%)	19	12	n.s.
Sleep attacks (%)	6.8	3.4	n.s.
Total sleep time (min)	450	442	n.s.
Snoring (%)	26	35	n.s.
Referred sleep apnea (%)	4.1	5.4	n.s.
NAP (%)	55	23	0.005
Duration ( min)	38.8	26.9	0.05